

---

**Nuclear transfer nTreg model reveals fate-determining TCR-beta and novel peripheral nTreg precursors.**

**Journal:** Proc Natl Acad Sci U S A

**Publication Year:** 2016

**Authors:** Manching Ku, Shih-En Chang, Julio Hernandez, Justin R Abadejos, Mohsen Sabouri-Ghomi, Niklas J Muenchmeier, Anna Schwarz, Anna M Valencia, Oktay Kirak

**PubMed link:** 27044095

**Funding Grants:** California State University-San Marcos CIRM Bridges to the Stem Cell Research Training Grant

**Public Summary:**

To study the development and function of "natural-arising" T regulatory (nTreg) cells, we developed a novel nTreg model on pure nonobese diabetic background using epigenetic reprogramming via somatic cell nuclear transfer. On RAG1-deficient background, we found that monoclonal FoxP3+CD4+Treg cells developed in the thymus in the absence of other T cells. Adoptive transfer experiments revealed that the thymic niche is not a limiting factor in nTreg development. In addition, we showed that the T-cell receptor (TCR) beta-chain of our nTreg model was not only sufficient to bias T-cell development toward the CD4 lineage, but we also demonstrated that this TCR beta-chain was able to provide stronger TCR signals. This TCR-beta-driven mechanism would thus unify former per se contradicting hypotheses of TCR-dependent and -independent nTreg development. Strikingly, peripheral FoxP3-CD4+T cells expressing the same TCR as this somatic cell nuclear transfer nTreg model had a reduced capability to differentiate into Th1 cells but were poised to differentiate better into induced nTreg cells, both in vitro and in vivo, representing a novel peripheral precursor subset of nTreg cells to which we refer to as pre-nTreg cells.

**Scientific Abstract:**

To study the development and function of "natural-arising" T regulatory (nTreg) cells, we developed a novel nTreg model on pure nonobese diabetic background using epigenetic reprogramming via somatic cell nuclear transfer. On RAG1-deficient background, we found that monoclonal FoxP3+CD4+Treg cells developed in the thymus in the absence of other T cells. Adoptive transfer experiments revealed that the thymic niche is not a limiting factor in nTreg development. In addition, we showed that the T-cell receptor (TCR) beta-chain of our nTreg model was not only sufficient to bias T-cell development toward the CD4 lineage, but we also demonstrated that this TCR beta-chain was able to provide stronger TCR signals. This TCR-beta-driven mechanism would thus unify former per se contradicting hypotheses of TCR-dependent and -independent nTreg development. Strikingly, peripheral FoxP3-CD4+T cells expressing the same TCR as this somatic cell nuclear transfer nTreg model had a reduced capability to differentiate into Th1 cells but were poised to differentiate better into induced nTreg cells, both in vitro and in vivo, representing a novel peripheral precursor subset of nTreg cells to which we refer to as pre-nTreg cells.

---

**Source URL:** <http://www.cirm.ca.gov/about-cirm/publications/nuclear-transfer-ntreg-model-reveals-fate-determining-tcr-beta-and-novel>